

2020-2021 Missouri School Immunization Requirements ~~~ Grades K-12

* There is a 4 day grace period for all types of vaccines

GR	DTaP/DTP/DT/Td/Tdap*	MENINGOCOCCAL*	POLIO*	MEASLES*	MUMPS*	RUBELLA*	HEPATITIS B*	VARICELLA*
	4+ DOSES		3+ DOSES	2 DOSES	2 DOSES	2 DOSES	3+ DOSES	2 DOSES
K-7	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses		Last dose on or after 4th birthday and at least 6 months after previous dose. If 4 doses before 4 years of age, a 5th dose should be given at 4 to 6 years of age.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Last dose must be administered on or after 168 days of age. Dose 1 - Initial visit. Dose 2 -28 days after Dose 1. Dose 3 - Minimum 56 days between dose 2 & 3. Cannot complete series in less than 4 months.	Dose 1 on or after first birthday and a minimum of 28 days between dose 1 and dose 2. (Recommended 3 months) or a statement signed by MD or DO indicating the month and year the child had chickenpox disease.
	4+ DOSES	1 DOSE	3+ DOSES	2 DOSES	2 DOSES	2 DOSES	3+ DOSES	2 DOSES
8-10	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses <u>1 Tdap required</u>	1 dose MCV required (must be given after age 10)	Last dose on or after fourth (4th) birthday. If any combination of 4 doses are administered appropriately and received prior to fourth birthday no additional dose is needed. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Dose 1 - Initial visit. Dose 2 -28 days after Dose 1. Dose 3 - Minimum 56 days between dose 2 & 3. Cannot complete series in less than 4 months.	Dose 1 on or after first birthday and a minimum of 28 days between dose 1 and dose 2. (Recommended 3 months) or a statement signed by MD or DO indicating the month and year the child had chickenpox disease..
	4+ DOSES	1 DOSE	3+ DOSES	2 DOSES	2 DOSES	2 DOSES	3+ DOSES	1 DOSE
11	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses <u>1 Tdap required</u>	1 dose MCV required (must be given after age 10)	Last dose on or after fourth (4th) birthday. If any combination of 4 doses are administered appropriately and received prior to fourth birthday no additional dose is needed. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Dose 1 - Initial visit. Dose 2 -28 days after Dose 1. Dose 3 - Minimum 56 days between dose 2 & 3. Cannot complete series in less than 4 months.	On or after first birthday or a statement signed by parent/guardian or physician indicating the month and year the child had chickenpox disease.
	4+ DOSES	2 DOSES	3+ DOSES	2 DOSES	2 DOSES	2 DOSES	3+ DOSES	1 DOSE
12	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses <u>1 Tdap required</u>	If first dose given at age 16 or older then only 1 dose is required.	Last dose on or after fourth (4th) birthday. If any combination of 4 doses are administered appropriately and received prior to fourth birthday no additional dose is needed. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Dose 1 - Initial visit. Dose 2 -28 days after Dose 1. Dose 3 - Minimum 56 days between dose 2 & 3. Cannot complete series in less than 4 months.	On or after first birthday or a statement signed by parent/guardian or physician indicating the month and year the child had chickenpox disease.